



Procedure Reminders

Colonoscopy/Flexible Sigmoidoscopy

Your Appointment

Appointment Date: _____ Arrival Time: _____

Your arrival time is 45 minutes prior to your procedure time. This allows time for check-in, health history review, and pre-procedure evaluation. Please be aware that we may contact you in the days leading up to your procedure, as your arrival time may be subject to change.

Reminders About Your Appointment

Please follow all instructions carefully, as improper preparation may result in your procedure being rescheduled.

- **Transportation:** Due to the sedation you will receive, someone at least 18 years of age MUST bring you to your appointment and drive you home afterwards. A taxi, bus, Uber, Lyft, or similar service is NOT an acceptable form of transportation. If you arrive without a driver, your procedure will be rescheduled. If you have questions about whether a method of transportation is acceptable, please contact our office at the number listed below.
- **Alcohol Consumption:** Do not consume any alcohol during the course of your procedure preparation.
- **Jewelry:** Please leave all jewelry and other valuables at home or with your driver, as we do not provide a secure area for these items.
- **Questions?** Contact our office at (336) 448-2427 (option 6, then option 2) if you have any questions about your prep or if there are any changes in your medical history between the day of scheduling and the appointment date.

Reminders About Medications

If you are taking medication on a regular basis:

- Take all necessary medications at least 4 hours prior to your arrival time.
- **Diabetes:** Please contact the prescribing physician regarding instructions for injectable diabetic medication the day before and the day of the procedure. DO NOT TAKE oral diabetic medication the day of the procedure.
- **Blood Pressure:** DO NOT SKIP any doses of your blood pressure medication.
- **Iron:** STOP taking iron 5 days before your procedure.
- **Inhalers and Nebulizers:** May be used up to the time of your procedure.
- **Blood Thinners:** Please contact the physician who prescribes this medication for you to make sure it is okay to stop taking it for the recommended time (listed below) prior to your procedure. You may remain on aspirin unless otherwise instructed. If you are on a blood thinner that is not listed, please contact the prescribing physician for instructions.

Aggrenox/Plavix: 5-7 days

Arixtra: 24 hours

Brilinta/Coumadin/Jantoven: 5 days

Effient: 7 days

Eliquis: 48 hours

Pletal/Persantine: 48 hours

Pradaxa: 48 hours

Savaysa: 1-3 days

Xarelto: 24-48 hours

FOLLOW THESE INSTRUCTIONS – NOT THOSE ON THE BOX YOU RECEIVE FROM THE PHARMACY!

Items to Purchase

Suprep (prescription)

2 Days Before Procedure


- ✓ Eat a well-balanced diet. Avoid whole grain foods, nuts, seeds, popcorn, raw or dried fruits and vegetables, and salads.
No solid food after midnight.

Day Before Procedure Beginning at 12:01am

- ✓ **Begin a clear liquid diet. DO NOT consume anything red in color OR dairy products.**
 Clear liquids may include water, soft drinks, coffee or tea without cream, popsicles, broth/bouillon, Jell-O, non-pulpy juice, etc.
- ✓ **At 4:00pm, complete steps 1 through 4 using only one of the 6-ounce bottles.**




Pour one 6-ounce bottle of Suprep liquid into the mixing container.



Add cool drinking water to the 16-ounce line on the container and mix.



Drink ALL the liquid in the container.



You **MUST** drink two more 16-ounce containers of water over the next hour.

Day of Procedure

- ✓ Continue a clear liquid diet.
- ✓ **6 hours prior to arrival time:** Repeat **steps 1-4** (shown above) using the other 6-ounce bottle of Suprep liquid.
- ✓ **4 hours prior to arrival time:** You must be finished with your final glass of water.
ABSOLUTELY NOTHING ELSE BY MOUTH AFTER THIS TIME!